arXiv:gr-qc/0612009v1 1 Dec 2006

Quasinormal modes of gravitational perturbation around a Schwarzschild black hole surrounded by quintessence

Yu Zhang* and Y.X.Gui†‡
Department of Physics, Dalian University of Technology, Dalian, 116024, People's Republic of China

In this paper, the quasinormal modes of gravitational perturbation around a Schwarzschild black hole surrounded by quintessence were evaluated by using the third-order WKB approximation. Due to the presence of quintessence, the gravitational wave damps more slowly.

PACS numbers: 02.60.-x, 04.30.-w, 95.30.Sf, 97.60.Lf
Keywords: Quasinormal modes; gravitational perturbation; WKB approximation.

I. INTRODUCTION

The quasinormal modes of a black hole present complex frequencies that correspond to solutions of the perturbation equations, which satisfy the boundary conditions appropriate for purely ingoing waves at the horizon and purely outgoing waves at infinity. Vishveshwara[1] firstly pointed out the concept of QNMs in calculations of the scattering of gravitational waves by a black hole. The study of QNMs of black hole has a long history[2]-[17]. The WKB[18]-[20] approach is a semi-analysis method which was used by many scientists to calculate QNMs such as Schwarzschild[21], Reissener-Norstrom[22], Kerr[23], Kerr-Newman[24]and so on . Recently, the properties of quasinormal modes have been investigated in the context of the AdS/CFT[25]-[31] correspondence and Loop Quantum Gravity[32]-[34].

Recent measurements of the cosmic microwave background(CMB)[35] combined with supernovae of type Ia[36], large scale structure (cosmic shear) [37] and galaxy cluster abundances[38] show that 70% of the Universe is dominated by a mysterious dark energy which causes the cosmic expansion to accelerate and 26% cold dark matter, the ordinary baryonic matter makes up only 4%. There are several types of models of dark energy, such as: the cosmological constant[39], quintessence[40], phantom[41], k-essence[42], quintom[43] models. For quintessence, the equation of state is given by $p_q = w_q\rho_q$ with w_q in the range of $-1 \leq w_q \leq -\frac{1}{3}$. Recently, Kiselev[44] considered Einstein's field equations for a black hole surrounded by the quintessential matter and obtained a new solution dependent on the state parameter w_q of the quintessence. And Songbai Chen et al[45] has evaluated the quasinormal frequencies of massless scalar field perturbation around the black hole which is surrounded by quintessence. The result shows that due to the presence of quintessence, the scalar field damps more rapidly. In our paper, we investigate gravitational perturbation in this situation.

In this paper, the quasinormal modes of Schwarzschild black hole surrounded by quintessence were evaluated by the third-order WKB approximation.

II. QUASINORMAL MODE OF GRAVITATIONAL PERTURBATION AROUND THE BLACK HOLE SURROUNDED BY QUINTESSENCE

The author[44] gets the general forms of exact spherically-symmetric solutions for the Einstein equations describing black holes surrounded by the quintessence with the energy momentum tensor, which satisfies the condition of the additivity and linearity. The metric[45] is given by

$$ds^2 = (1 - \frac{2M}{r} - \frac{c}{r^{3w_q+1}})dt^2 - (1 - \frac{2M}{r} - \frac{c}{r^{3w_q+1}})^{-1}dr^2 - r^2(d\theta^2 + sin\theta^2 d\phi^2), \tag{1}$$

where M is the black hole mass, w_q is the quintessential state parameter, c is the normalization factor related to $\rho_q = -\frac{c}{2}\frac{3w_q}{r^{3(1+w_q)}}$, and ρ_q is the density of quitenssence.

The study of black hole perturbations was initiated by Regge and Wheeler[46] for the ‘odd ’parity types of harmonics and was continued to the ‘even’parity by Zerilli[47].

---

† Corresponding author
*Electronic address: zhangyu128@student.dlut.edu.cn
‡Electronic address: thphys@dlut.edu.cn

Indicate the background metric with $g_{\mu\nu}$ and the perurbation in it with $h_{\mu\nu}$. The perturbation $h_{\mu\nu}$ are supposed to be very small compared with $g_{\mu\nu}$. The $h_{\mu\nu}$ can be calculated from $g_{\mu\nu}$, and $R_{\mu\nu}+\delta R_{\mu\nu}$ from $g_{\mu\nu}+h_{\mu\nu}$. $\delta R_{\mu\nu}$ can be expressed in the form[48]

$$\delta R_{\mu\nu} = -\delta\Gamma^{\beta}_{\mu\nu;\,\beta} + \delta\Gamma^{\beta}_{\mu\beta;\,\nu}, \tag{2}$$

where

$$\delta\Gamma^{\alpha}_{\beta\gamma} = \frac{1}{2} g^{\alpha\nu}(h_{\beta\nu;\,\gamma} + h_{\gamma\nu;\,\beta} - h_{\beta\gamma;\,\nu}), \tag{3}$$

The canonical form for the perturbations in the Regge-Wheeler gauge is given as[46]

$$h_{\mu\nu} = \begin{vmatrix} 0 & 0 & 0 & h_0(r) \\ 0 & 0 & 0 & h_1(r) \\ 0 & 0 & 0 & 0 \\ sym & sym & 0 & 0 \end{vmatrix} \exp(-i\omega T)(\sin\theta/\partial\theta)P_L(\cos\theta) \tag{4}$$

Submit (4) to (2), we can get

$$(1-\frac{2M}{r}-\frac{c}{r^{3w_q+1}})^{-1} i\omega h_0 + \frac{d}{dr}[(1-\frac{2M}{r}-\frac{c}{r^{3w_q+1}})h_1] = 0 \qquad from \quad \delta R_{23}=0 \tag{5}$$

$$(1-\frac{2M}{r}-\frac{c}{r^{3w_q+1}})^{-1} i\omega(\frac{dh_0}{dr} + i\omega h_1 - \frac{2h_0}{r}) + (l-1)(l+2)\frac{h_1}{r^2} = 0 \qquad from \quad \delta R_{13}=0 \tag{6}$$

Defining

$$\Phi(r) = (1-\frac{2M}{r}-\frac{c}{r^{3w_q+1}})h_1/r \tag{7}$$

and

$$\frac{dr_*}{dr} = (1-\frac{2M}{r}-\frac{c}{r^{3w_q+1}})^{-1} \tag{8}$$

and eliminate h_0, we then get

$$(\frac{d^2}{dr_*^2}+\omega^2)\Phi(r) = V\Phi(r) \tag{9}$$

where

$$V = (1-\frac{2M}{r}-\frac{c}{r^{3w_q+1}})(\frac{l(l+1)}{r^2} - \frac{6M}{r^3} - \frac{c(3w_q+3)}{r^{3w_q+3}}) \tag{10}$$

In this paper, we use the third-order WKB approximation method devised by Schutz, Will[18] and Iyer[19].The formula for the complex quasinormal frequencies ω is

$$\omega^2 = \left[V_0 + (-2V_0'')^{1/2}\Lambda\right] - i(n+\frac{1}{2})(-2V_0'')^{1/2}(1+\Omega) \tag{11}$$

where

$$\Lambda = \frac{1}{(-2V_0'')^{1/2}}\left\{\frac{1}{8}\left(\frac{V_0^{(4)}}{V_0''}\right)(\frac{1}{4}+\alpha^2) - \frac{1}{288}\left(\frac{V_0'''}{V_0''}\right)^2(7+60\alpha^2)\right\}$$

$$\begin{aligned}\Omega = \frac{1}{(-2V_0'')}\Big\{&\frac{5}{6912}\left(\frac{V_0'''}{V_0''}\right)^4(77+188\alpha^2) \\ &- \frac{1}{384}\left(\frac{V_0'''^2 V_0^{(4)}}{V_0''^3}\right)(51+100\alpha^2) + \frac{1}{2304}\left(\frac{V_0^{(4)}}{V_0''}\right)^2(67+68\alpha^2) \\ &+ \frac{1}{288}\left(\frac{V_0''' V_0^{(5)}}{V_0''^2}\right)(19+28\alpha^2) - \frac{1}{288}\left(\frac{V_0^{(6)}}{V_0''}\right)(5+4\alpha^2)\Big\}\end{aligned} \tag{12}$$

and

$$\alpha = n + \frac{1}{2},\ V_0^{(n)} = \frac{d^n V}{dr_*^n}\Big|_{r_*=r_*(r_p)} \tag{13}$$

Take $M = 1, c = 0.001$ and $M = 1, c = 0$ for our calculation. And $c = 0$ means there is no quintessence. Using the third-order WKB approximation, we can get the solutions as shown in the table 1 and table 2 . Where l is the angular harmonic index, n is the overtone number, ω is the complex quasinormal frequencie, w_q is the quintessential state parameter.

TABLE I: The quasinormal frequencies of gravitational perturbation in a Schwarzshild black hole without quintessence(c=0).

l	n	ω	l	n	ω
2	0	$0.37316 - 0.08922i$	4	0	$0.80910 - 0.09417i$
	1	$0.34602 - 0.27492i$		1	$0.79650 - 0.28437i$
	2	$0.30293 - 0.47106i$		2	$0.77364 - 0.47897i$
	3	$0.24746 - 0.67290i$		3	$0.74331 - 0.67830i$
				4	$0.70721 - 0.88127i$
3	0	$0.59927 - 0.09273i$	5	0	$1.01225 - 0.09487i$
	1	$0.58235 - 0.28141i$		1	$1.00215 - 0.28583i$
	2	$0.55320 - 0.47668i$		2	$0.98326 - 0.47990i$
	3	$0.51575 - 0.67743i$		3	$0.95748 - 0.67780i$
	4	$0.47107 - 0.88154i$		4	$0.92636 - 0.87919i$

III. DISCUSSION AND CONCLUSION

The data of table I is obtained by using the WKB method in a Schwarzschild black hole without quintessence, and table II is under the quintessence. Explicitly, we plot the relationship between the real and imaginary parts of quasinormal frequencies with the variation of w_q(for fixed $c = 0.001$), and compared with no quintessence. From the figure I we can find that for fixed c(unequal to 0) and l the absolute value of the real and imaginary parts decrease as the quintessence state parameter w_q decreases. The quasi-normal modes of the black hole present complex frequencies whose real part represents the actual frequency of the oscillation and the imaginary part representing the damping. It means that when the value of w_q is smaller, the oscillations damps more slowly. Also the absolute value of the real and imaginary parts of quasinormal modes with quintessence is smaller compared with no quintessence for given l and n. That is to say, due to the presence of quintessence, the oscillations of the gravitational wave damps more slowly, and that is different from the situation of the scalar field given by Songbai Chen et al[45].

As we know, lots of stars will end their lives with a violent supernova explosion. This will leave behind a compact object oscillated violently in the first few seconds, emit a large amount of gravitational radiation and the initial oscillations will damp out. We could observe for a few tenths of a millisecond continuous monochromatic blasting if the supernova remnants formed a black hole. And the gravitational waves will carry away information about the black hole. From the calculation, the paper shows that the quintessence will influence the quasinormal modes of gravitational perturbation around a black hole. We set $c = 0.001$. c is the normalization factor related to the density of quintessence around the black hole. Actually, c may be much smaller than 0.001 and hence the influence to gravitational perturbation could be neglected. Although the QNMs are predicted by the perturbation equations, it is not always clear which ones will be excited and under what initial conditions. Only when the density of the quintessence surrounding the black hole is high enough to make the influence on the QNMs distinct and measurable could we investigate the character of quintessence by way of extraction and analysing the experimental data.

Acknowledgments

Yu Zhang wishes to thank Professor H Y Liu and Dr LiXin Xu for helpful discussions. This work is supported by the National Natural Science Foundation of China under Grant No. 10573004.

TABlE II:The quasinormal frequencies of gravitational perturbation in a Schwarzshild black hole surrounded by quintessence for $l = 2$, $l = 3$, $l = 4$, $l = 5$ and $c = 0.001$.

$3w_q + 1$	$\omega(n = 0)$	$\omega(n = 1)$	$\omega(n = 2)$	$\omega(n = 3)$	$\omega(n = 4)$
$l = 2$					
0	$0.37261 - 0.08904i$	$0.34553 - 0.27437i$	$0.30256 - 0.47013i$	$0.24722 - 0.67155i$	
-0.4	$0.37230 - 0.08894i$	$0.34527 - 0.27406i$	$0.30235 - 0.46958i$	$0.24710 - 0.67078i$	
-0.8	$0.37182 - 0.08880i$	$0.34485 - 0.27362i$	$0.30205 - 0.46882i$	$0.24693 - 0.66967i$	
-1.2	$0.37107 - 0.08862i$	$0.34423 - 0.27303i$	$0.30161 - 0.46776i$	$0.24672 - 0.66810i$	
-1.6	$0.36990 - 0.08840i$	$0.34333 - 0.27227i$	$0.30105 - 0.46628i$	$0.24651 - 0.66585i$	
-2.0	$0.36810 - 0.08819i$	$0.34209 - 0.27131i$	$0.30039 - 0.46415i$	$0.24629 - 0.66240i$	
$l = 3$					
0	$0.59837 - 0.09255i$	$0.58151 - 0.28085i$	$0.55242 - 0.47574i$	$0.51506 - 0.67608i$	$0.47048 - 0.87978i$
-0.4	$0.59788 - 0.09244i$	$0.58104 - 0.28052i$	$0.55199 - 0.47518i$	$0.51468 - 0.67528i$	$0.47017 - 0.87874i$
-0.8	$0.59712 - 0.09229i$	$0.58031 - 0.28006i$	$0.55134 - 0.47438i$	$0.51410 - 0.67415i$	$0.46968 - 0.87726i$
-1.2	$0.59592 - 0.09208i$	$0.57919 - 0.27942i$	$0.55033 - 0.47329i$	$0.51324 - 0.67257i$	$0.46899 - 0.87518i$
-1.6	$0.59406 - 0.09183i$	$0.57748 - 0.27862i$	$0.54887 - 0.47184i$	$0.51204 - 0.67039i$	$0.46807 - 0.87223i$
-2.0	$0.59117 - 0.09156i$	$0.57495 - 0.27767i$	$0.54684 - 0.46992i$	$0.51049 - 0.66729i$	$0.46690 - 0.86787i$
$l = 4$					
0	$0.80789 - 0.09398i$	$0.79532 - 0.28380i$	$0.77252 - 0.47802i$	$0.74226 - 0.67694i$	$0.70625 - 0.87950i$
-0.4	$0.80723 - 0.09387i$	$0.79468 - 0.28347i$	$0.77190 - 0.47745i$	$0.74169 - 0.67614i$	$0.70572 - 0.87846i$
-0.8	$0.80619 - 0.09372i$	$0.79367 - 0.28299i$	$0.77094 - 0.47664i$	$0.74079 - 0.67499i$	$0.70490 - 0.87696i$
-1.2	$0.80457 - 0.09350i$	$0.79210 - 0.28234i$	$0.76947 - 0.47554i$	$0.73943 - 0.67340i$	$0.70367 - 0.87487i$
-1.6	$0.80206 - 0.09324i$	$0.78970 - 0.28151i$	$0.76726 - 0.47409i$	$0.73746 - 0.67126i$	$0.70195 - 0.87199i$
-2.0	$0.79815 - 0.09294i$	$0.78606 - 0.28055i$	$0.76406 - 0.47228i$	$0.73474 - 0.66840i$	$0.69966 - 0.86795i$
$l = 5$					
0	$1.01074 - 0.09468i$	$1.00066 - 0.28526i$	$0.98182 - 0.47894i$	$0.95610 - 0.67644i$	$0.92505 - 0.87743i$
-0.4	$1.00990 - 0.09457i$	$0.99984 - 0.28492i$	$0.98103 - 0.47837i$	$0.95534 - 0.67563i$	$0.92433 - 0.87638i$
-0.8	$1.00861 - 0.09441i$	$0.99856 - 0.28444i$	$0.97979 - 0.47756i$	$0.95415 - 0.67448i$	$0.92321 - 0.87488i$
-1.2	$1.00658 - 0.09419i$	$0.99658 - 0.28378i$	$0.97788 - 0.47644i$	$0.95234 - 0.67289i$	$0.92151 - 0.87280i$
-1.6	$1.00344 - 0.09392i$	$0.99353 - 0.28294i$	$0.97499 - 0.47499i$	$0.94966 - 0.67078i$	$0.91906 - 0.86999i$
-2.0	$0.99854 - 0.09362i$	$0.98884 - 0.28198i$	$0.97068 - 0.47324i$	$0.94581 - 0.66808i$	$0.91567 - 0.86619i$

[1] C. V. Vishveshwara, Nature **227** 936 (1970).

[2] S. Chandrasekhar, Proc. R. Soc. London, Ser. A. **343**, 289-298 (1975).; S. Chandrasekhar, and S. Detweiler, Proc. R. Soc. London, Ser. A. **344**, 441-452 (1975).

[3] H. P. Nollert, Phys. Rev. D **47**, 5253-5258 (1993).

[4] E. W. Leaver, Proc. R. Soc. A **402** 285 (1985) .

[5] K. D. Kokkotas and B. G. Schmidt, Living Rev. Rel. **2**2 (1999); H. P. Nollert, Class. Quant. Grav. **16**, R159 (1999).

[6] V. Cardoso and J. P. S. Lemos Phys. Rev. D 64, 084017 (2001); E. Beri, V. Cardoso and S. Yoshida, Phys. Rev. D69 124018 (2004); V. Cardoso, J. P. S. Lemos, S. Yoshida, Phys. Rev. D**70** 124032 (2004) .

[7] A. J. M. Medved, D. Martin, M. Visser, Class. Quant. Grav. **21** 2393-2405 (2004) . A. J. M. Medved, D. Martin, Gen. Rel. Grav. **37** 1529-1539 (2005)

[8] H. B. Zhang, Z. J. Cao, X. F. Gong, W. Zhou,Class. Quant. Grav. **21** 917-925 (2004)

[9] J. L. Jing, Phys. Rev. D**69** 084009 (2004) ; S. B. Chen, J. L. Jing, Class. Quant. Grav. **22** 533-539 (2005) ; S. B. Chen, J. L. Jing, Class. Quant. Grav. **22** 2159-2165 (2005) .

[10] R. Konoplya, Phys. Rev. D**71** 024038 (2005).

[11] I. B. Khriplovich, Int. J. Mod. Phys. D**14** 181 (2005).

[12] F. W. Shu, Y.G. Shen, Phys. Rev. D**70** 084046 (2004).

[13] J. Crisóstomo, S. Lepe, J. Saavedra, Class. Quant. Grav. **21** 2801-2809 (2004).

[14] F. W. Shu, Y. G. Shen, Phys. Lett. B**614** 195-200 (2005).
[15] R. G. Daghigh, G. Kunstatter, Class. Quant. Grav. **22** 4113-4128 (2005).
[16] E. Berti, K. D. Kokkotas, Phys. Rev. D**71** 124008 (2005).
[17] M. Giammatteo, I. G. Moss, Class. Quant. Grav. **22** 1803-1824 (2005).
[18] B. F. Schutz, and C. M. Will, Astrophys. J.**291**, L33-L36 (1985).
[19] S. Iyer and C. M. Will, Phys. Rev. D **35**, 3621 (1987)
[20] R. A. Konoplya, Phys. Rev. D **68** 024018 (2003)
[21] S. Iyer, Phys. Rev. D **35**, 3632 (1987);
[22] K. D. Kokkotas and B. F. Schutz, Phys. Rev. D **37**, 3378 (1988)
[23] E. Seidel, and S. Iyer, Phys. Rev. D **41**, 374-382 (1990).
[24] K. D. Kokkotas, Nuovo Cimento B **108**, 991-998 (1993).
[25] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 1998.
[26] E. Witten, Adv. Theor. Math. Phys. **2**, 253 1998.
[27] Gubser S S Phys. Rev. D**63** 084017 2001
[28] B. Wang, C. Y. Lin and E. Abdalla, Phys. Lett. B **481**, 79 (2000). B. Wang, E. Abdalla and R. B. Mann, Phys. Rev. D **65**, 084006 (2002); B. Wang, C.Y. Lin and C. Molina, Phys. Rev. D **70**, 064025 (2004).
[29] V. Cardoso and J. P. S. Lemos, Phys. Rev. D **64**, 084017 (2001); V. Cardoso and J. P. S. Lemos, Class. Quantum. Grav. **18**, 5257 (2001).
[30] D. Birmingham, I. Sachs, S. N. Solodukhin, Phys. Rev. Lett. **88**, 151301 (2002); D. Birmingham, Phys. Rev. D **64**, 064024 (2001).
[31] R. A. Konoplya, Phys. Rev. D **66**, 044009 (2002).
[32] S. Hod, Phy. Rev. Lett. **81**, 4293 (1998).
[33] A. Corichi, Phys. Rev. D **67**, 087502 (2003).
[34] O. Dreyer, Phy. Rev. Lett. **90**, 081301 (2003).
[35] A. D. Miller et al., Astrophys. J. Lett. **524**, L1 (1999); P. de Bernardis et al., Nature **404**, 955 (2000); S. Hanany et al., Astrophys. J. Lett. **545**, L5 (2000); N. W. Halverson et al., Astrophys. J. **568**, 38 (2002); L. Page et al., Astrophys. J. Suppl. **148**, 233 (2003); S. Hannestad, G. Raffelt, Phys. Rev. D **72** 103514 (2005); R. Lamon, R. Durrer, Phys. Rev. D **73** 023507 (2006).
[36] P. Ruiz-Lapuente, A. Burkert, R. Canal, Astrophys. J. **447** L69 (1995); A. G. Riess et al., Astron. J. **116**, 1009 (1998); D. Branch, Ann. Rev. Astron.Astrophys. **36** 17-55 (1998); R. A. Knop et al., Astrophys. J. **598**, 102 (2003). A. G. Riess et al., Astrophys. J. **607**, 665 (2004); X. Zhang, F. Q. Wu, Phys. Rev. D **72** 043524 (2005).
[37] D. J. Bacon et al., MNRAS, **318**, 625 (2000). D. J. Bacon et al., MNRAS, **344**, 673 (2003). R. Scranton et al., astro-ph/0307335; N. Kaiser et al., astro-ph/0003338; M. Tegmark et al., Phys. Rev. D **69**, 103501 (2004).
[38] T. Kitayama, Y. Suto, Astrophys. J. **469** 480 (1996) ; J. V. Cunha, J. S. Alcaniz, J. A. S. Lima, Phys. Rev. D**69** 083501 (2004); Y. Wang, M. Tegmark, Phys. Rev. Lett. **92** 241302 (2004) .
[39] S. Weinberg, Rev. Mod. Phys. **61**, 1 (1989); K. A. Olive, M. Pospelov, Phys. Rev. D **65** 085044 (2002); J. S. Alcaniz, Phys. Rev. D **69** 083521 (2004).
[40] B. Ratra and P. J. E. Peebles, Phys. Rev. D **37**, 3406 (1988); P. J. E. Peebles and B. Ratra, Astrophys. J. **325**, L17 (1988); R. RCaldwell, R. Dave, and P. J. Steinhardt, Phys. Rev. Lett. **80**, 1582 (1998); I. Zlatev, L. Wang and P. J. Steinhardt, Phys. Rev. Lett. **82**, 896 (1999); V. Sahni, L. M. Wang, Phys. Rev. D **63**, 103517 (2000); T. Matos, L. A. Urena-Lopez , Phys. Rev. D **63**, 063506 (2001); S. Capozziello, V.F. Cardone, E. Piedipalumbo, C. Rubano, Class.Quant.Grav. **23**, 1205-1216 (2006).
[41] A. E. Schulz, M. White, Phys. Rev. D **64**, 043514 (2001); R. R. Caldwell, Phys. Lett. B **545**, 23 (2002); S. Nojiri and S. D. Odintsov, Phys. Lett. B **562**, 147 (2003); S. Nojiri and S. D. Odintsov, Phys. Lett. B **565**, 1 (2003); P. Singh, M. Sami, and N. Dadhich, Phys. Rev. D **68**, 023522 (2003); L. P. Chimento and R. Lazkoz, Phys. Rev. Lett. **91**, 211301 (2003); A. Vikman, Phys. Rev. D **71**, 023515 (2005).
[42] T. Chiba, T. Okabe and M. Yamaguchi, Phys. Rev. D **62**, 023511 (2000); R. J. Scherrer, Phys. Rev. Lett. **93**, 011301 (2004); L. P. Chimento, M. Forte, R. Lazkoz, Mod. Phys. Lett. A **20** 2075 (2005).
[43] B. Feng, M. Z. Li, Y. S. Piao, X. M. Zhang, Phys. Lett. B **634** 101-105 (2006) ; H. Wei, R. G. Cai, D. F. Zeng, Class. Quant. Grav. **22** 3189-3202 (2005); G. B. Zhao, J. Q. Xia, M. Z. Li, B. Feng, X. M. Zhang, Phys. Rev. D **72** 123515(2005).
[44] V. V. Kiselev, Class. Quant. Grav. **20**, 1187-1197 (2003).
[45] S. B. Chen, J. L. Jing, Class. Quant. Grav. **22**, 4651-4657 (2005).
[46] T. Regge , J. A. Wheeler, Phys. Rev., V. **108**, 1063-1069 (1957).
[47] F. J. Zerilli, Phys. Rev. Lett. **24**, 737 (1970).
[48] L. P. Einsenhart, Riemannian Geometry, Princeton Univ, press, Princeton (1926), Chap. VI.



FIG. 1: The relationship between the real and imaginary parts of quasinormal frequencies of the gravitational perturbation in the background of the black hole surrounded by quintessence for fixed $c = 0.001$ and no quintessence for $c = 0$. $\times$ and $*$ refer to that there is quintessence and no quintessence, respectively.